# Multi-criteria analysis on agriculture exposure to drought in arid areas

ESCAP developed a methodology to assess the agriculture land exposure to drought-related climate risk using aridity and projections for environmental risk. The agriculture exposure to multi-criteria indicators for drought was modified based on the Land suitability assessment method for agriculture in arid land developed by Marieme Seif-Ennasr and others (2020). There are 3 main variables used in this analysis, namely 1) Aridity, 2) Climate-related data, 3) Slope, and 4) Agriculture data as exposure.

Once the aridity, climate and slope indicators were spatially analyzed through scoring and normalization; these were then combined. Afterwards, the output was overlaid with agriculture exposure and the exposure of agriculture to certain level of drought risks was quantified.

Multi-criteria analysis on Agriculture exposed to drought in arid areas (aridity, decrease of rainfall, increasing number of dry days and temperature and slope)

Agriculture land

#### **Aridity**

\*

Normalized score of aridity for arid & semi arid (Low score of aridity → more arid)



#### Drought

Normalized % Change in Maximum 5-day precipitation amount during dry days (in June-August)

Threshold → only the negative percentages

Higher precipitation → less drought-risk possibility

#### Temperature change

Normalized Increasing Temperature **change** (unit: deg C). Threshold >0.

USAID, 2018: Increased temperature change → more damage to crops in Central Asia

### Number of dry days

Normalized Increasing Number of consecutive dry day s (unit: days, pr<1mm). Threshold >0



**Slope** from DEM (0.1, 0.3, 0.6, and 1)

Seif-Ennasr et al, 2020: degree slope → better land suitability → less risk

ESCAP modified from "Land suitability assessment for agriculture method" by Ma rieme Seif-Ennasr et al, 2020.

## The following diagram shows the indicators and datasets used in our analysis.

